

This Page Is Inserted by IFW Operations  
and is not a part of the Official Record

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- ✓ • BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning documents *will not* correct images,  
please do not report the images to the  
Image Problem Mailbox.**

## FIGURES

Figure 1A (schematic drawing)

### ZsProSensor-1





Figure 1B: sequence


```
atggcccagtccaagcacggcctgaccaaggagatgacatgaagtaccgcatggagggctgcgtggatggccaca  
gttcgtgatcacggcgagggcatcggtacccctcaagggcaagcaggccatcaacctgtgcgtggtggagggcgg  
cccttgcccttgcgcgaggacatcttgccgcgccttcaactacggcaaccgcgtgttcaccgagtaccccaggacat  
cgtcgactactcaagaactcctgccccgcggctacacctgggaccgctccttctgttcgaggacggcgccgtgtgcat  
ctgcaacgccgacatcacctgagcgtggaggagaactgcatgtaccacgagtcgaagtctacggcgtgaactcccc  
gccgacggccccgtgatgaagaagatgaccgacaactgggagccctctgcgagaagatcatccccgtgccaagca  
gggcatcttgaagggcgacgtgagcatgtacctgctgctgaaggacggtggccgcttgccgtgccagttcgacaccgtg  
acaaggccaagtccgtgccccgcaagatgcccgactggcacttcattccagcacaagctgacccgcgaggaccgcagcg  
acgccaagaaccagaagtggcacctgaccgagcacgccatcgctccggtccgccttgccccgcggtcacggccaa  
tgtggcaactcatgaaacagatccagagccatggcttcccgcggaggtggaggagcaggatgatggcacgctgccca  
tgtcttgctcccaggagagcgggatggaccgtcacctgcagcctgtgcttctgctaggatcaatgtg
```

Figure 2: Targeting of ZsGreen to degradation by the proteasome using motif from MODC.

**Figure 2A:** Flow Cytometry. Mean Fluorescence Intensities (MFI) of HEK 293 cells transiently transfected with plasmids encoding ZsGreen, ZsGreend1 and ZsGreend410. Standard deviations from duplicates.




**Figure 2B:** Flow Cytometry. Same as 1A. Cells were treated for 0 to 6 hours with 10 ug/ml ALLN.



**Figure 3: Generation of stable cell clone expressing ZsGreend410 to monitor the activity of the proteasome in a HTS fashion.**

**Figure 3A: Flow Cytometry.** MFI of a stable clone of HEK 293 transfected with a plasmid encoding ZsGreend410. Cells treated for 6 hours with or without 10 ug/ml ALLN. Standard deviations from duplicates.



**Figure 3B: Microscopy.** Micrographs of a stable clone of HEK 293 transfected with a plasmid encoding ZsGreend410. Cells treated for 10 hours with 10 ug/ml ALLN. Micrographs taken with same exposure times.

**Figure 3C: 96 well plate fluorescence reader.** Standard deviation from triplicates.




FIGURE 4 (same as 2B with the stable clone)






Figure 5A, B, C and D. Dose response curve obtained with the stable clone and Acumen explorer machine. Compound 1=Epoxomycin; compound 2=Lactacystin; compound 3=ZLLH; compound 4=ALLN.


Compound One - Proteosome Sensor Assay (650Volts PMT Settings) - Peak Intensity



Compound Two - Proteosome Sensor Assay (650Volts PMT Settings) - Peak Intensity



**Compound Three - Proteosome Sensor Assay (650Volts PMT Settings) - Peak Intensity**



**Compound Four - Proteosome Sensor Assay (650Volts PMT Settings) - Peak Intensity**

